

# An Insight into Trigonometric Identities and Trigonometric Equations

## Elementary Trigonometric Results

Recall definition of sine, cosine and tangent of an angle.

Let  $P(x, y)$  be a point on a circle with radius  $r = \sqrt{x^2 + y^2}$  in the Cartesian Plane


As usual,  $\angle XOP$  is measured in the direction from  $\overline{OX}$  to  $\overline{OP}$ , taking anticlockwise as positive.

Let  $\theta = \angle XOP$

$$\text{Then } \sin \theta = \frac{y}{r}, \quad \cos \theta = \frac{x}{r}, \quad \tan \theta = \frac{\sin \theta}{\cos \theta} = \frac{y}{x}$$

## Relations between trigonometric functions of angles of opposite signs

Observe the following diagram where points A and A', B and B' are reflection of each other in the x-axis (think of A' to A as well as A to A')



We can summarize the change of coordinates corresponding to change of sign of angle (namely a reflection) as this :

$$\begin{pmatrix} x \\ y \end{pmatrix} \rightarrow \begin{pmatrix} x \\ -y \end{pmatrix} \text{ as } \theta \rightarrow -\theta, \text{ for all values of } \theta, \text{ positive or negative}$$


We thus have the result

$$\sin(-\theta) = -\sin \theta, \quad \cos(-\theta) = \cos \theta \text{ and } \tan(-\theta) = -\tan \theta \dots\dots\dots (1)$$

## Relations between trigonometric functions of angles by a difference of one right angle

The following diagram shows how coordinates are changed as a result of adding one right angle to a given one. The various cases are:

$\angle XOP$  is changed to  $\angle XOQ$ ,  $\angle XOQ$  is changed to  $\angle XOR$ , .....



We can summarize the change of coordinates as a result of the addition of one right angle as this:

$$\begin{pmatrix} x \\ y \end{pmatrix} \rightarrow \begin{pmatrix} -y \\ x \end{pmatrix} \text{ as } \theta \rightarrow \theta + \frac{\pi}{2}, \text{ where } \theta \text{ is in any quadrant}$$

We thus have

$$\text{If } \cos \theta = \frac{x}{r} \text{ and } \sin \theta = \frac{y}{r},$$

$$\text{then } \sin\left(\theta + \frac{\pi}{2}\right) = \frac{x}{r} = \cos \theta, \quad \cos\left(\theta + \frac{\pi}{2}\right) = \frac{-y}{r} = -\sin \theta$$

So,

$$\sin\left(\theta + \frac{\pi}{2}\right) = \cos \theta, \quad \cos\left(\theta + \frac{\pi}{2}\right) = -\sin \theta \text{ and } \tan\left(\theta + \frac{\pi}{2}\right) = -\tan \theta$$

$$\text{for all values of } \theta \dots\dots\dots (2)$$

(The reader should not just take  $\angle XOP$  as an example of  $\theta$ , instead, think about  $\angle XOQ$ ,  $\angle XOR$  and  $\angle XOS$  as well)


With results (1) and (2), which are valid for all values of  $\theta$ , we deduce all results related to complimentary and supplementary angles.

For example,

$$\cos(180^\circ - \theta) = \cos(90^\circ + 90^\circ - \theta) = -\sin(90^\circ - \theta) = -\cos(-\theta) = -\cos \theta$$

which is true for all values of  $\theta$

**Formulae for addition and subtraction of angles.**



The above figure shows two angles  $p$  and  $q$  of any size, given by  $\angle AOC$  and  $\angle AOB$  respectively, where A, B and C are points on a circle, centre O, radius  $r$ . (It need not be  $p > q$  as depicted)

Let OA be lying on the OX axis and so the angles are measured using OX as initial direction.

We find  $C(r \cos p, r \sin p)$ ,  $B(r \cos q, r \sin q)$

By using the cosine rule, we have, for all possible positions of B and C,

$$2(OB)(OC)\cos(p - q) = (OB)^2 + (OC)^2 - (BC)^2$$

So,

$$\begin{aligned} 2r^2 \cos(p - q) &= r^2 + r^2 - [(r \cos p - r \cos q)^2 + (r \sin p - r \sin q)^2] \\ &= 2r^2 - r^2[\cos^2 p - 2 \cos p \cos q + \cos^2 q] - r^2[\sin^2 p - 2 \sin p \sin q + \sin^2 q] \\ &= 2r^2 - r^2[1 + 1 - 2 \cos p \cos q - 2 \sin p \sin q] \\ &= 2r^2[\cos p \cos q + \sin p \sin q] \end{aligned}$$

$$\text{So, } \cos(p - q) = \cos p \cos q + \sin p \sin q, \text{ for all values of } p \text{ and } q \dots\dots\dots (3)$$

From results (1), (2) and (3) we can establish all formulae related to additions and subtractions of angles,

For example

$$\cos(p + q) = \cos(p - (-q)) = \cos p \cos -q + \sin p \sin -q = \cos p \cos q - \sin p \sin q$$

### Trigonometric Equations

Let us investigate the three simple equations

$$\sin \theta = a, \text{ where } -1 \leq a \leq 1$$

$$\cos \theta = b, \text{ where } -1 \leq b \leq 1$$

$$\tan \theta = c, \text{ for all values of } c$$

First note that all three equations have two solutions in any interval of length  $2\pi$ , for example, we can find two solutions in each of  $(0.5\pi, 2.5\pi)$ ,  $(-0.9\pi, 1.1\pi)$  etc  
 So, to begin with, it is usually convenient to obtain two in the range of  $[-\pi, \pi]$

Next, we note that if  $\alpha$  is a solution, say to  $\sin \theta = a$ , where  $-1 \leq a \leq 1$ , then  $\sin \alpha = a$ ,  $\sin(\alpha + 2\pi) = a$ ,  $\sin(\alpha - 2\pi) = a$ ,  $\sin(\alpha + 2n\pi) = a$ , where  $n$  is any integer  
 So, if  $\alpha, \beta$  are two solutions to  $\sin \theta = a$ , where  $-1 \leq a \leq 1$ , in the interval  $[-\pi, \pi]$ , then the complete solution set is  $\theta = \alpha + 2n\pi, \beta + 2n\pi$ , though of course, it works out to be  $\theta = n\pi + (-1)^n(\alpha)$ , which is equivalent to  $\theta = n\pi + (-1)^n(\beta)$   
 (More discussions are left as an exercise)

**Exercise (without answers attached)**

Recall these results:

$$\sin(-\theta) = -\sin \theta, \cos(-\theta) = \cos \theta \text{ and } \tan(-\theta) = -\tan \theta, \text{ for all values of } \theta \dots\dots\dots (1)$$

$$\sin(\theta + \frac{\pi}{2}) = \cos \theta, \cos(\theta + \frac{\pi}{2}) = -\sin \theta \text{ and } \tan(\theta + \frac{\pi}{2}) = -\tan \theta \text{ for all values of } \theta \dots\dots\dots (2)$$

$$\cos(p - q) = \cos p \cos q + \sin p \sin q, \text{ for all values of } p \text{ and } q \dots\dots\dots (3)$$

1. Testify the rule :  $\begin{pmatrix} x \\ y \end{pmatrix} \rightarrow \begin{pmatrix} -y \\ x \end{pmatrix}$  as  $\theta \rightarrow \theta + 90^\circ$ , where P(x, y) is the point on a circle of radius  $r$ ,  $\theta = \angle XOP$ , for the following cases
2. By using Results (1) and (2) above, express the following in terms of  $\sin \theta, \cos \theta$  and  $\tan \theta$  only
  - (i)  $\sin(180^\circ - \theta)$  (ii)  $\cos(180^\circ - \theta)$  (iii)  $\tan(180^\circ - \theta)$
  - (iv)  $\sin(180^\circ + \theta)$  (v)  $\cos(180^\circ + \theta)$  (vi)  $\tan(180^\circ + \theta)$
3. By using Results (1) and (2) above, express the following in terms of  $\sin \theta, \cos \theta, \tan \theta$  and  $\cot \theta$  only
  - (i)  $\sin(270^\circ - \theta)$  (ii)  $\cos(270^\circ - \theta)$  (iii)  $\tan(270^\circ - \theta)$
  - (iv)  $\sin(270^\circ + \theta)$  (v)  $\cos(270^\circ + \theta)$  (vi)  $\tan(270^\circ + \theta)$
4. By using Results (1), (2) and (3) above, obtain the formulae  $\sin(p + q) = \sin p \cos q + \cos p \sin q$  and  $\sin(p - q) = \sin p \cos q - \cos p \sin q$

5.  $\alpha, \beta$  are two different solutions of the equation  
 $\sin \theta = a$ , where  $|a| \leq 1$   
State a relation between  $\alpha$  and  $\beta$  if  
(i)  $\alpha$  and  $\beta$  are in the same quadrant  
(ii)  $\alpha$  and  $\beta$  are in different quadrant  
(iii) there is no information about the quadrants of  $\alpha$  and  $\beta$ .
6.  $\alpha, \beta$  are two different solutions of the equation  
 $\cos \theta = a$ , where  $|a| \leq 1$   
State a relation between  $\alpha$  and  $\beta$
7.  $\alpha, \beta$  are two different solutions of the equation  
 $\tan \theta = a$   
What is the values of  $\alpha - \beta$  ?
8. In the solution for the equation  $\cos(2x - 70^\circ) = 0.5$ , values of  $x$  required for  
are in the interval  $(-150^\circ, 570^\circ)$   
How many possible values of  $x$  are there?

**Exercise (with answers attached)**

Recall these results:

$$\sin(-\theta) = -\sin \theta, \cos(-\theta) = \cos \theta \text{ and } \tan(-\theta) = -\tan \theta, \\ \text{for all values of } \theta \dots\dots\dots (1)$$

$$\sin(\theta + \frac{\pi}{2}) = \cos \theta, \cos(\theta + \frac{\pi}{2}) = -\sin \theta \text{ and } \tan(\theta + \frac{\pi}{2}) = -\tan \theta \\ \text{for all values of } \theta \dots\dots\dots (2)$$

$$\cos(p - q) = \cos p \cos q + \sin p \sin q, \text{ for all values of } p \text{ and } q \dots\dots\dots (3)$$

1. Testify the rule :

$\begin{pmatrix} x \\ y \end{pmatrix} \rightarrow \begin{pmatrix} -y \\ x \end{pmatrix}$  as  $\theta \rightarrow \theta + 90^\circ$ , where P(x, y) is the point on a circle of radius r,  $\theta = \angle XOP$ , for the following cases

- (i)  $\theta = 240^\circ$       (ii)  $\theta = -240^\circ$       (iii)  $\theta = 500^\circ$

2. By using Results (1) and (2) above, express the following in terms of  $\sin \theta$ ,  $\cos \theta$  and  $\tan \theta$  only

- (i)  $\sin(180^\circ - \theta)$     (ii)  $\cos(180^\circ - \theta)$     (iii)  $\tan(180^\circ - \theta)$   
 (iv)  $\sin(180^\circ + \theta)$     (v)  $\cos(180^\circ + \theta)$     (vi)  $\tan(180^\circ + \theta)$

Answers

- (i)  $\sin \theta$               (ii)  $-\cos(\theta)$           (iii)  $-\tan \theta$   
 (iv)  $-\sin \theta$           (v)  $-\cos \theta$             (vi)  $\tan \theta$

3. By using Results (1) and (2) above, express the following in terms of  $\sin \theta$ ,  $\cos \theta$ ,  $\tan \theta$  and  $\cot \theta$  only

- (i)  $\sin(270^\circ - \theta)$     (ii)  $\cos(270^\circ - \theta)$     (iii)  $\tan(270^\circ - \theta)$   
 (iv)  $\sin(270^\circ + \theta)$     (v)  $\cos(270^\circ + \theta)$     (vi)  $\tan(270^\circ + \theta)$

Answers:

- (i)  $-\cos \theta$             (ii)  $-\sin \theta$             (iii)  $\cot \theta$   
 (iv)  $-\cos \theta$           (v)  $\sin \theta$               (vi)  $-\cot \theta$

4. By using Results (1), (2) and (3) above, obtain the formulae

$$\sin(p + q) = \sin p \cos q + \cos p \sin q \text{ and} \\ \sin(p - q) = \sin p \cos q - \cos p \sin q$$

Solution

$$\sin(p + q) = -\cos(\frac{\pi}{2} + p + q) = -[\cos(\frac{\pi}{2} + p)\cos q - \sin(\frac{\pi}{2} + p)\sin q] \\ = \sin p \cos q + \cos p \sin q$$

$$\sin(p - q) = -\cos(\frac{\pi}{2} + p - q) = -[\cos(\frac{\pi}{2} + p)\cos -q - \sin(\frac{\pi}{2} + p)\sin -q]$$

$$= \sin p \cos q - \cos p \sin q$$

5.  $\alpha, \beta$  are two different solutions of the equation  
 $\sin \theta = a$ , where  $|a| \leq 1$

State a relation between  $\alpha$  and  $\beta$  if

- (i)  $\alpha$  and  $\beta$  are in the same quadrant
- (ii)  $\alpha$  and  $\beta$  are in different quadrant
- (iv) there is no information about the quadrants of  $\alpha$  and  $\beta$ .

Answers:

- (i)  $\beta = 2n\pi + \alpha$ , where  $n \in \mathbb{Z}$
- (ii)  $\beta = 2n\pi + \pi - \alpha$ , where  $n \in \mathbb{Z}$   $\pi - \alpha$
- (iii)  $\beta = \pi - \alpha, 2\pi + \alpha, 3\pi - \alpha, \dots, -\pi - \alpha, -2\pi + \alpha, -3\pi - \alpha, \dots$   
 So,  $\beta = n\pi + (-1)^n \alpha$ , where  $n \in \mathbb{Z}$

6.  $\alpha, \beta$  are two different solutions of the equation  
 $\cos \theta = a$ , where  $|a| \leq 1$

State a relation between  $\alpha$  and  $\beta$

Answer

$$\beta = 2n\pi \pm \alpha, \text{ where } n \in \mathbb{Z}$$

7.  $\alpha, \beta$  are two different solutions of the equation  
 $\tan \theta = a$

What is the values of  $\alpha - \beta$  ?

Answer

$$\alpha - \beta = n\pi, \text{ for } n \in \mathbb{Z}$$

- 8 In the solution for the equation  $\cos(2x - 70^\circ) = 0.5$ , values of  $x$  required for are in the interval  $(-150^\circ, 570^\circ)$

How many possible values of  $x$  are there?

Answer

$$\text{The interval length for } x \text{ is } 570^\circ - (-150^\circ) = 720^\circ = 2 \times 360^\circ$$

$$\text{So, the interval length for } (2x - 70^\circ) \text{ is } 4 \times 360^\circ$$

So, there are 8 possible values for  $(2x - 70^\circ)$ , and therefore also 8 possible values for  $x$ .