


This article was downloaded by: [Dr Weng Kin Ho]

On: 21 December 2011, At: 21:29

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



International Journal of Mathematical Education in Science and Technology

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/tmes20>

An elementary proof of the identity

Weng Kin Ho^a, Foo Him Ho^a & Tuo Yeong Lee^b

^a National Institute of Education, Singapore 637616, Singapore

^b NUS High School of Math & Science, Singapore 129957, Singapore

Available online: 21 Dec 2011

To cite this article: Weng Kin Ho, Foo Him Ho & Tuo Yeong Lee (2011): An elementary proof of the identity, International Journal of Mathematical Education in Science and Technology, DOI:10.1080/0020739X.2011.644337

To link to this article: <http://dx.doi.org/10.1080/0020739X.2011.644337>



PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.tandfonline.com/page/terms-and-conditions>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

CLASSROOM NOTE

An elementary proof of the identity $\cot \theta = \frac{1}{\theta} + \sum_{k=1}^{\infty} \frac{2\theta}{\theta^2 - k^2\pi^2}$

Weng Kin Ho^{a*}, Foo Him Ho^a and Tuo Yeong Lee^b

^aNational Institute of Education, Singapore 637616, Singapore

^bNUS High School of Math & Science, Singapore 129957, Singapore

(Received 25 July 2011)

This article gives an elementary proof of the famous identity

$$\cot \theta = \frac{1}{\theta} + \sum_{k=1}^{\infty} \frac{2\theta}{\theta^2 - k^2\pi^2}, \quad \theta \in \mathbb{R} \setminus \pi\mathbb{Z}.$$

Using nothing more than freshman calculus, the present proof is far simpler than many existing ones. This result also leads directly to Euler's and Neville's identities, as well as the identity $\zeta(2) := \sum_{k=1}^{\infty} \frac{1}{k^2} = \frac{\pi^2}{6}$.

Keywords: Euler's identity; partial fractions; calculus

1. Introduction

The famous identity,

$$\pi \cot(\pi z) = \frac{1}{z} + \sum_{v=1}^{\infty} \frac{2z}{z^2 - v^2}, \quad z \in \mathbb{C} \setminus \mathbb{Z} \quad (1)$$

was probably known to Euler, who later in 1770 presented a more general (and more complicated) form

$$\frac{\pi}{n} \cdot \frac{\cos \frac{\pi(w-z)}{2n}}{\sin \frac{\pi(w+z)}{2n} - \sin \frac{\pi(w-z)}{2n}} = \frac{1}{z} + \sum_{v=1}^{\infty} \left(\frac{2w}{(2v-1)^2 n^2 - w^2} - \frac{2z}{(2v)^2 n^2 - z^2} \right)$$

in [1], where $n \in \mathbb{Z}^+$ and $w, z \in \mathbb{C}$. Equation (1) which is also known as the *partial fraction representation* of $\pi \cot(\pi z)$ can be shown to hold for non-integer complex numbers z , using complex analysis [2–4]. In particular, [4] is the first to put in print a proof that employs the so-called *Herglotz's trick*, thereby simplifying the proof given in [3]. For an excellent historical account of this identity and more, the reader is referred to [5].

The real version of (1), expressed as

$$\cot x = \frac{1}{x} + \sum_{k=1}^{\infty} \frac{2x}{x^2 - k^2\pi^2}, \quad x \in \mathbb{R} \setminus \pi\mathbb{Z}, \quad (2)$$

*Corresponding author. Email: wengkin.ho@nie.edu.sg

is already an interesting identity in itself, attracting numerous elegant proofs using real analysis [6,7]. Unfortunately, many of these proofs are quite involved and thus not readily accessible by beginners in calculus.

This article presents an elementary and simpler proof of (2), using no more than freshman calculus. Because of its lighter mathematical overhead, it is hoped that the material herein can be of educational value in the following sense. Instructors of calculus courses can make suitable adaptations of this material for classroom teaching and discussion, design of investigative activities and enrichment experiences. This article is also written with teacher training purposes in mind, i.e., to supply trainee teachers with pedagogical content knowledge relevant to the teaching of calculus.

We organize this article as follows. In Section 2, we establish some elementary inequalities essential in the proof of the main result. Our proof of Equation (2) appears in Section 3. The section is written in a self-contained manner, i.e., each of the results we employ can be proven with background knowledge acquired from a first-year undergraduate Calculus course. In doing so, we demonstrate the usefulness of the crucial results commonly taught in such a course. We intentionally and completely bypass *convergence theorems*; whenever an interchange of limits, infinite sums and (differential or integral) operators is invoked, there is always an elementary alternative. Section 4 showcases some interesting applications of the identity (2). This serves resource materials that can be employed in the design of classroom activities for a freshman course in Calculus. Throughout this article, we denote the set of integers, integral multiples of π , real numbers and complex numbers by \mathbb{Z} , $\pi\mathbb{Z}$, \mathbb{R} and \mathbb{C} , respectively.

2. Some inequalities

We start with two basic inequalities commonly encountered in Calculus.

Proposition 2.1: *If $t \in (0, \pi)$, then $t - \frac{t^3}{3!} \leq \sin t \leq t - \frac{t^3}{3!} + \frac{t^5}{5!}$.*

Proof: We establish only the lower bound, leaving the upper one as an exercise for the diligent reader. Consider $g(t) = \sin t - (t - \frac{t^3}{3!})$. Since $g''(t) = -\sin t + t \geq 0$, it follows from $g'(0) = 0$ that $g'(t) = \cos t - (1 - \frac{t^2}{2!}) \geq 0$. Because $g(0) = 0$, it then follows that $g(t) \geq 0$. \square

Proposition 2.2: *If $t \in (0, \frac{\pi}{2})$, then $\frac{t}{\sin t} \leq \frac{\pi}{2}$.*

Proof: This follows since $f(t) = \frac{\sin t}{t}$ is monotone decreasing on $(0, \frac{\pi}{2})$. \square

These results lead to the following handy inequality:

Lemma 2.3: *If $t \in (0, \frac{\pi}{2})$, then $0 < -\frac{1}{t^2} + \csc^2 t < 1$.*

Proof: Since $0 < \sin t < t$ for all $t \in (0, \frac{\pi}{2})$, we have $0 < \frac{1}{t^2} < \csc^2 t$ so that $0 < -\frac{1}{t^2} + \csc^2 t$. Now, by Proposition 2.2,

$$-\frac{1}{t^2} + \csc^2 t = \frac{(t - \sin t)(t + \sin t)}{t^2 \sin^2 t} \leq \frac{(t - t + \frac{t^3}{3!})(2t)}{t^2 \sin^2 t} = \frac{1}{3} \left(\frac{t}{\sin t} \right)^2 \leq \frac{1}{3} \cdot \frac{\pi^2}{4} < 1.$$

\square

3. Partial fraction representation of $\cot \theta$

The topic of partial fractions is key to many national curricula for pre-university or high school mathematics (e.g. [8,9]). Apart from its usual role in series expansion and antiderivatives of rational functions, its connection with other elementary functions is seldom mentioned.

In this section, our main aim is to obtain the following partial fraction representation for $\cot \theta$:

Theorem 3.1: For $\theta \in \mathbb{R} \setminus \pi\mathbb{Z}$, it holds that

$$\cot \theta = \frac{1}{\theta} + \sum_{k=1}^{\infty} \frac{2\theta}{\theta^2 - k^2\pi^2}. \quad (3)$$

Such a partial fraction representation must necessarily involve an infinite sum since any finite partial fraction amounts to only a rational function, *not* a trigonometric one. Because an infinite series can be seen as the limit of a recursion, it seems fit to initiate our proof from a common-looking trigonometric identity which somewhat ‘recursively’ defines $\cot \theta$.

Proposition 3.2: For any $x \in (0, \pi)$, the following identity holds:

$$\cot x = \frac{1}{2} \left(\cot \frac{x}{2} - \tan \frac{x}{2} \right).$$

Proof: Take reciprocals in the double angle formula $\tan 2x = \frac{2 \tan x}{1 - \tan^2 x}$. □

With a recursive mind-set, we apply the preceding proposition thrice to obtain

$$\begin{aligned} \cot \theta &= \frac{1}{2} \left(\cot \frac{\theta}{2} - \tan \frac{\theta}{2} \right) \\ &= \frac{1}{2} \left(\frac{1}{2} \cot \frac{\theta}{4} - \frac{1}{2} \tan \frac{\theta}{4} - \tan \frac{\theta}{2} \right) \\ &= \frac{1}{2^2} \left(\cot \frac{\theta}{2^2} - \tan \frac{\theta}{2^2} \right) - \frac{1}{2} \cot \left(\frac{\pi}{2} - \frac{\theta}{2} \right) \\ &= \frac{1}{2^2} \left(\cot \frac{\theta}{2^2} - \tan \frac{\theta}{2^2} \right) - \frac{1}{2} \cdot \frac{1}{2} \left(\cot \frac{\pi - \theta}{2^2} - \cot \frac{\pi + \theta}{2^2} \right) \\ &= \frac{1}{2^2} \left(\cot \frac{\theta}{2^2} - \tan \frac{\theta}{2^2} \right) + \frac{1}{2^2} \left(\cot \frac{\theta + \pi}{2^2} + \cot \frac{\theta - \pi}{2^2} \right). \end{aligned}$$

A straightforward induction on $n \geq 2$ then yields the generalization below.

Proposition 3.3: If $\theta \in (0, \pi)$ and $n \geq 2$ is an integer, then

$$\cot \theta = \frac{1}{2^n} \left(\cot \frac{\theta}{2^n} - \tan \frac{\theta}{2^n} \right) + \frac{1}{2^n} \sum_{k=1}^{2^{n-1}-1} \left(\cot \frac{\theta + k\pi}{2^n} + \cot \frac{\theta - k\pi}{2^n} \right).$$

The well-worn fact that $\lim_{\alpha \rightarrow 0} \frac{\alpha}{\sin \alpha} = 1$ justifies the following proposition.

Proposition 3.4: If $\theta \in (0, \pi)$, then

$$\lim_{n \rightarrow \infty} \frac{1}{2^n} \left(\cot \frac{\theta}{2^n} - \tan \frac{\theta}{2^n} \right) = \frac{1}{\theta}.$$

Based on Propositions 3.3 and 3.4, it becomes evident that to achieve (3) one must prove the following theorem.

Theorem 3.5: *If $\theta \in (0, \pi)$, then*

$$\lim_{n \rightarrow \infty} \left\{ \frac{1}{2^n} \sum_{k=1}^{2^{n-1}-1} \left(\cot \frac{\theta + k\pi}{2^n} + \cot \frac{\theta - k\pi}{2^n} \right) - \sum_{k=1}^{2^{n-1}-1} \frac{2\theta}{\theta^2 - k^2\pi^2} \right\} = 0.$$

Proof: For $n \geq 2$, we have:

$$\begin{aligned} & \frac{1}{2^n} \sum_{k=1}^{2^{n-1}-1} \left(\cot \frac{\theta + k\pi}{2^n} + \cot \frac{\theta - k\pi}{2^n} \right) - \sum_{k=1}^{2^{n-1}-1} \frac{2\theta}{\theta^2 - k^2\pi^2} \\ &= \frac{1}{2^n} \sum_{k=1}^{2^{n-1}-1} \left(\cot \frac{\theta + k\pi}{2^n} - \frac{2^n}{\theta + k\pi} + \cot \frac{\theta - k\pi}{2^n} - \frac{2^n}{\theta - k\pi} \right). \\ &= \frac{1}{2^n} \sum_{k=1}^{2^{n-1}-1} \int_{\frac{k\pi-\theta}{2^n}}^{\frac{k\pi+\theta}{2^n}} \left(\frac{1}{t^2} - \csc^2 t \right) dt. \end{aligned}$$

Since $0 < -\frac{1}{t^2} + \csc^2 t \leq 1$ for all $t \in \left(\frac{k\pi-\theta}{2^n}, \frac{k\pi+\theta}{2^n} \right) \subset (0, \frac{\pi}{2})$ by Lemma 2.3,

$$0 \leq \frac{1}{2^n} \sum_{k=1}^{2^{n-1}-1} \int_{\frac{k\pi-\theta}{2^n}}^{\frac{k\pi+\theta}{2^n}} \left(-\frac{1}{t^2} + \csc^2 t \right) dt \leq \frac{\theta}{2^n}.$$

By the Squeeze theorem, it follows that

$$\lim_{n \rightarrow \infty} \frac{1}{2^n} \sum_{k=1}^{2^{n-1}-1} \int_{\frac{k\pi-\theta}{2^n}}^{\frac{k\pi+\theta}{2^n}} \left(\frac{1}{t^2} - \csc^2 t \right) dt = 0$$

and consequently,

$$\lim_{n \rightarrow \infty} \left\{ \frac{1}{2^n} \sum_{k=1}^{2^{n-1}-1} \left(\cot \frac{\theta + k\pi}{2^n} + \cot \frac{\theta - k\pi}{2^n} \right) - \sum_{k=1}^{2^{n-1}-1} \frac{2\theta}{\theta^2 - k^2\pi^2} \right\} = 0.$$

□

All in all, we have succeeded in showing the following lemma.

Lemma 3.6: *If $\theta \in (0, \pi)$, the following identity holds:*

$$\cot \theta = \frac{1}{\theta} + \sum_{k=1}^{\infty} \frac{2\theta}{\theta^2 - k^2\pi^2}.$$

To prove Theorem 3.1, we need to extend Lemma 3.6 for $\theta \in \mathbb{R} \setminus \pi\mathbb{Z}$. To prove this assertion, we choose $\alpha \in (0, \pi)$ and $r \in \mathbb{Z}$ so that $\theta = \alpha + r\pi$.

Since $\cot(\theta - r\pi) = \cot \theta$, the result follows from Lemma 3.6 that

$$\begin{aligned} \cot \theta &= \cot(\theta - r\pi) \\ &= \frac{1}{\theta - r\pi} + \sum_{k=1}^{\infty} \frac{2(\theta - r\pi)}{(\theta - r\pi - k\pi)(\theta - r\pi + k\pi)} \end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{\theta - r\pi} + \sum_{k=1}^{\infty} \left(\frac{1}{\theta - r\pi - k\pi} + \frac{1}{\theta - r\pi + k\pi} \right) \\
 &= \frac{1}{\theta - r\pi} + \sum_{k=r+1}^{\infty} \left(\frac{1}{\theta - k\pi} + \frac{1}{\theta + (k-r)\pi} \right) + \sum_{k=1}^r \frac{1}{\theta + (k-r)\pi} \\
 &= \frac{1}{\theta - r\pi} + \sum_{k=r+1}^{\infty} \left(\frac{1}{\theta - k\pi} + \frac{1}{\theta + (k-r)\pi} \right) + \sum_{k=0}^{r-1} \frac{1}{\theta - k\pi} \\
 &= \frac{1}{\theta} + \sum_{k=1}^{\infty} \left(\frac{1}{\theta - k\pi} + \frac{1}{\theta + k\pi} \right) \\
 &= \frac{1}{\theta} + \sum_{k=1}^{\infty} \frac{2\theta}{\theta^2 - k^2\pi^2}.
 \end{aligned}$$

The proof of Theorem 3.1 is now complete.

4. Some applications

Putting $\theta = \pi z$ into (3), $z \notin \mathbb{Z}$, yields immediately:

Corollary 4.1 (Euler's identity): *For $z \in \mathbb{R} \setminus \mathbb{Z}$, it holds that*

$$\pi \cot(\pi z) = \frac{1}{z} + \sum_{k=1}^{\infty} \frac{2z}{z^2 - k^2}.$$

An interesting consequence of Theorem 3.1, which is almost immediate, is:

Corollary 4.2 (Neville's identity): *If $\theta \in \mathbb{R} \setminus \pi\mathbb{Z}$, then*

$$\csc^2 \theta - \frac{1}{\theta^2} = \sum_{k=1}^{\infty} \left(\frac{1}{(\theta - k\pi)^2} + \frac{1}{(\theta + k\pi)^2} \right). \quad (4)$$

A naïve derivation of (4) is via a term-by-term differentiation of (3) since $\frac{2\theta}{\theta^2 - k^2\pi^2} = \frac{1}{\theta - k\pi} + \frac{1}{\theta + k\pi}$. However, this approach involves the unjustified interchange of the differential operator and the infinite sum. We present below a proof to fully justify this interchange of operators without appealing to any convergence theorems.

Proof: Choose $\delta\theta > 0$ so small that $(\theta - \delta\theta, \theta + \delta\theta) \subset (r\pi, (r+1)\pi)$ for some $r \in \mathbb{Z}$. Then for any integer $n > |r| + 1$, we have

$$\begin{aligned}
 &\frac{1}{\delta\theta} \left| \sum_{k=1}^n \int_{\theta}^{\theta+\delta\theta} \left(\frac{1}{(t - k\pi)^2} + \frac{1}{(t + k\pi)^2} \right) dt - \int_{\theta}^{\theta+\delta\theta} \sum_{k=1}^{\infty} \left(\frac{1}{(t - k\pi)^2} + \frac{1}{(t + k\pi)^2} \right) dt \right| \\
 &= \frac{1}{\delta\theta} \left| \int_{\theta}^{\theta+\delta\theta} \sum_{k=n+1}^{\infty} \left(\frac{1}{(t - k\pi)^2} + \frac{1}{(t + k\pi)^2} \right) dt \right| \\
 &\leq \frac{1}{\pi^2 \delta\theta} \left| \int_{\theta}^{\theta+\delta\theta} \sum_{k=n+1}^{\infty} \left(\frac{1}{(k - r - 1)^2} + \frac{1}{(k + r)^2} \right) dt \right|
 \end{aligned}$$

$$\begin{aligned}
&< \sum_{k=n+1}^{\infty} \left(\frac{1}{(k-r-1)^2} + \frac{1}{(k+r)^2} \right) \\
&< \frac{2}{n-1-r}.
\end{aligned}$$

Letting $n \rightarrow \infty$, we have:

$$\sum_{k=1}^{\infty} \int_{\theta}^{\theta+\delta\theta} \left(\frac{1}{(t-k\pi)^2} + \frac{1}{(t+k\pi)^2} \right) dt = \int_{\theta}^{\theta+\delta\theta} \sum_{k=1}^{\infty} \left(\frac{1}{(t-k\pi)^2} + \frac{1}{(t+k\pi)^2} \right) dt.$$

Now, Theorem 3.1 asserts that

$$\frac{1}{\delta\theta} \left\{ \frac{1}{\theta + \delta\theta} - \cot(\theta + \delta\theta) - \left(\frac{1}{\theta} - \cot \theta \right) \right\} = \frac{1}{\delta\theta} \int_{\theta}^{\theta+\delta\theta} \sum_{k=1}^{\infty} \left(\frac{1}{(t-k\pi)^2} + \frac{1}{(t+k\pi)^2} \right) dt.$$

Taking $\delta\theta \rightarrow 0$, it follows that

$$\sum_{k=1}^{\infty} \left(\frac{1}{(\theta - k\pi)^2} + \frac{1}{(\theta + k\pi)^2} \right) = \csc^2 \theta - \frac{1}{\theta^2}$$

and the proof is now complete. \square

Remark 4.3: The above proof of (4) is much more direct than the original one presented in [6].

Corollary 4.4 (Euler's formula for $\zeta(2)$):

$$\zeta(2) := \sum_{k=1}^{\infty} \frac{1}{k^2} = \frac{\pi^2}{6}.$$

Proof: On the one hand,

$$\lim_{\theta \rightarrow 0^+} \sum_{k=1}^{\infty} \left(\frac{1}{(\theta - k\pi)^2} + \frac{1}{(\theta + k\pi)^2} \right) = \frac{2}{\pi^2} \sum_{k=1}^{\infty} \frac{1}{k^2}.$$

To account for the legitimate interchange of limits, we first observe, for $\theta \in (0, \frac{\pi}{2})$, that

$$\sum_{k=1}^{\infty} \frac{1}{(k\pi + \theta)^2} \leq \sum_{k=1}^{\infty} \frac{1}{k^2 \pi^2} \leq \sum_{k=1}^{\infty} \frac{1}{(k\pi - \theta)^2}.$$

Next, we need to establish that $\sum_{k=1}^{\infty} \left\{ \frac{1}{(k\pi - \theta)^2} - \frac{1}{(k\pi + \theta)^2} \right\} \leq 16\theta$ for $\theta \in (0, \frac{\pi}{2})$. This derives easily from the following inequalities:

$$\begin{aligned}
\sum_{k=1}^{\infty} \left\{ \frac{1}{(k\pi - \theta)^2} - \frac{1}{(k\pi + \theta)^2} \right\} &\leq \sum_{k=1}^{\infty} \frac{4k\pi\theta}{(k^2\pi^2 - \theta^2)^2} \\
&\leq \sum_{k=1}^{\infty} \frac{4k\theta}{\pi^3(k^2 - \frac{1}{4})^2} \\
&\leq \sum_{k=1}^{\infty} \frac{64k\theta}{(4k^2 - 1)^2} \leq \sum_{k=1}^{\infty} \frac{32\theta}{4k^2 - 1} = 16\theta,
\end{aligned}$$

where the third-to-second inequality holds since $4k^2 - 1 = (2k - 1)(2k + 1) \geq 2k$. Thus,

$$\lim_{\theta \rightarrow 0^+} \sum_{k=1}^{\infty} \frac{1}{(k\pi + \theta)^2} = \lim_{\theta \rightarrow 0^+} \sum_{k=1}^{\infty} \frac{1}{(k\pi - \theta)^2} = \sum_{k=1}^{\infty} \frac{1}{k^2 \pi^2}.$$

This then implies that

$$\lim_{\theta \rightarrow 0^+} \sum_{k=1}^{\infty} \left(\frac{1}{(\theta - k\pi)^2} + \frac{1}{(\theta + k\pi)^2} \right) = \frac{2}{\pi^2} \sum_{k=1}^{\infty} \frac{1}{k^2}.$$

On the other hand, Proposition 2.1 bears upon us to have

$$\frac{(\frac{\theta^3}{3!} - \frac{\theta^5}{5!})(2\theta - \frac{\theta^3}{3!})}{\theta^4} \leq \frac{(\theta - \sin \theta)(\theta + \sin \theta)}{\theta^4} \leq \frac{\frac{\theta^3}{3!} \cdot 2\theta}{\theta^4}$$

and thus, $\lim_{\theta \rightarrow 0^+} (\csc^2 \theta - \frac{1}{\theta^2}) = \frac{1}{3}$. The desired result then follows. \square

5. Concluding remarks

This article presents a simple and direct proof of the famous Euler's partial fraction representation of the cotangent function. Besides being an intriguing identity in its own right, (3) has seen recent applications in the parametric Euler's iterated sums [10]. We hope to demonstrate, within this small exposition, that even basic calculus techniques can be harnessed to yield non-trivial results that are usually obtained through, otherwise, more difficult means.

Acknowledgements

We are grateful to the anonymous referee for giving us invaluable advice that greatly improved our previous version of this article.

References

- [1] L. Euler, *De seriebus quibusdam considerationes*, Opera Omnia 1(14) (1970), pp. 407–462.
- [2] H. Schröter, *Ableitung der partialbruch- und produkt-entwickelungen für die trigonoem-trischen funktione*, Zeitschr. Math. Phys. 13 (1868), pp. 254–259.
- [3] F.H. Schottky, *Über das additiontheorem der cotangente*, J. Reine Angew. Math. 110 (1892), pp. 324–337.
- [4] C. Carathéodory, *Theory of Functions of a Complex Variable I*, F. Steinhart, trans., 2nd ed., Chelsea, New York, 1958 (in German).
- [5] R. Remmert, *Theory of Complex Functions*, Springer, New York, 1991.
- [6] E.H. Neville, *A trigonometrical inequality*, Proc. Camb. Philos. Soc. 47 (1951), pp. 629–632.
- [7] K. Knopp, *Theory and Application of Infinite Series*, Dover, New York, 1990.
- [8] Singapore Ministry of Education, *Additional Mathematics GCE 'O' Level Syllabus 4038*, 2011.
- [9] P.T. Joshi, *Efficient techniques for partial fractions*, Two-Year Coll. Math. J. 14(2) (1983), pp. 110–118.
- [10] D. Borwein, J.M. Borwein, and D.M. Bradley, *Parametric Euler sum identities*, J. Math. Anal. Appl. 316(1) (2004), pp. 328–338.